



ORIGINAL ARTICLE



Analysis of the impact of health interventions on vaccination coverage for children under two years of age in municipalities of Minas Gerais

Análise do impacto das intervenções em saúde sobre a cobertura vacinal para crianças menores de dois anos em municípios de Minas Gerais

Carolina Machado Moreira^I , Thales Philippe Rodrigues da Silva^{II,III} , Mariana Coelho De Almeida Neves^{IV} , Marcus Vinicius Gonçalves da Cruz^V , Elice Eliane Nobre Ribeiro^V , Silvio Ferreira Júnior^V , Sheila Aparecida Ferreira Lachtim^I , Fernanda Penido Matozinhos^I

^IUniversidade Federal de Minas Gerais – Belo Horizonte (MG), Brazil.

^{II}Universidade Federal de São Paulo, Escola Paulista de Enfermagem, Department of Women's Health Nursing – São Paulo (SP), Brazil.

^{III}Universidade Federal de Minas Gerais, School of Nursing, Postgraduate Program in Nursing – Belo Horizonte (MG), Brazil.

^{IV}Faculdade de Ciências Médicas de Minas Gerais – Belo Horizonte (MG), Brazil.

^VFundação João Pinheiro – Belo Horizonte (MG), Brazil.

ABSTRACT

Objective: To evaluate the impact of the state action-research project on vaccination coverage in children under two years of age in the state of Minas Gerais, according to the size of the municipalities, comparing the years 2021 and 2022. **Methods:** This is a study nested within the state action-research project, a before-after community clinical trial carried out in 212 municipalities in the state of Minas Gerais. This study used secondary data on Vaccination Coverage (VC), Homogeneity of Vaccines (HVC) and Abandonment rate of multi-dose vaccines. After classifying municipalities by size and vaccination coverage rates were equitably classified, an analysis of secondary data on 12 immunobiologicals indicated for the age group in question and their abandonment rate of multi-dose vaccines was carried out. **Results:** There was an increase in the proportion of municipalities classified as small that reached the vaccination coverage target set by the National Immunization Program (PNI) after the action-research project was carried out. There was an increase in the proportion of small municipalities classified as having a low abandonment rate for the rotavirus vaccine, in the adequate homogeneity of vaccination coverage and in the classification of risk as very low risk and low and medium risk, all with a statistically significant difference. **Conclusion:** There was an influence of municipal size on the effectiveness of the actions applied to increase vaccination coverage, explaining that proposing individualized actions for each municipality is essential to improve vaccination coverage.

Keywords: Vaccination. Vaccination coverage. Communicable diseases. Counties. Epidemiology.

CORRESPONDING AUTHOR: Fernanda Penido Matozinhos. Avenida Professor Alfredo Balena, 190, Santa Efigênia, CEP: 30130-100, Belo Horizonte (MG), Brasil. E-mail: nandapenido@hotmail.com

CONFLICT OF INTERESTS: nothing to declare

HOW TO CITE THIS ARTICLE: Moreira CM, Silva TPR, Neves MCA, Cruz MVG, Ribeiro EEN, Ferreira Júnior S, et al. Analysis of the impact of health interventions on vaccination coverage for children under two years of age in municipalities of Minas Gerais. Rev Bras Epidemiol. 2024; 27: e240028. <https://doi.org/10.1590/1980-549720240028>

This is an open article distributed under the CC-BY 4.0 license, which allows copying and redistribution of the material in any format and for any purpose as long as the original authorship and publication credits are maintained.

Received on: 09/25/2023

Reviewed on: 02/12/2024

Accepted on: 02/16/2024



INTRODUCTION

After five decades since the establishment of the National Immunization Program (*Programa Nacional de Imunizações* – PNI), Brazil remains a global benchmark in vaccination efforts, offering 19 immunization agents tailored for various life stages and specific populations. Through comprehensive coordination strategies, PNI has successfully eliminated and eradicated vaccine-preventable diseases nationwide^{1,2}, thereby ensuring effective protection for individuals and communities alike, while significantly enhancing the public health landscape.

Moreover, the implementation of PNI has focused on inclusivity and the reduction of regional and social disparities by operating in hard-to-reach areas without discrimination against individuals^{1,2}. This underscores the practical application of universality, a cornerstone principle of the Brazilian Unified Health System (*Sistema Único de Saúde* – SUS), which has transformed healthcare into a fundamental right for all^{1,2}.

However, it should be noted that the average vaccination coverage in Brazil declined from 97% in 2015 to 75% in 2020 for vaccines such as BCG, Hepatitis B, Poliomyelitis, Human Rotavirus, Pentavalent, Pneumococcal 10 and 13, Meningococcal C, Yellow Fever, and Triple Viral³. This decline has led to the concerning resurgence of diseases that were previously eradicated⁴⁻⁶.

This decline in vaccination coverage in recent years cannot be attributed to a single reason, as the drop is the result of multiple factors⁷⁻⁹. These include a lack of awareness about the seriousness of the situation, insufficient investments in health, structural issues and the circulation of fake news are reasons that possibly work together for the drop in the vaccination rate⁸. It is imperative to implement measures to restore high levels of vaccination coverage in the country^{10,11}.

Brazil includes states with vast territorial extensions, such as Minas Gerais, which comprises 853 municipalities, making it the state with the largest number of municipalities in the country. It ranks fourth in terms of territorial area, encompassing extensive urban and rural areas¹². The decline in vaccination coverage in Minas Gerais has mirrored similar trends observed in other states, particularly concerning the BCG, Polio, and MMR vaccines¹². Given its vast territorial expanse, Minas Gerais exhibits diverse environmental, cultural, and primarily social contexts. Hence, it is crucial to examine regional and quantitative variations in immunization across each municipality to establish a stratified overview, facilitating the formulation of targeted and uniform interventions^{11,12}.

The decline in vaccination coverage among children in Minas Gerais exhibited regional disparities in 2021: the state had 70.5% of municipalities experiencing very low Homogeneity of Vaccination Coverage (HVC). Interestingly, among these municipalities, those with larger populations

demonstrated the highest percentages of low coverage¹¹. In view of this, health regionalization must act to reduce these disparities and this risk factor for public health in the country and so that there are no setbacks in the achievements of PNI^{1,4,13}.

The objective of the study conducted by the Center for Studies and Research in Vaccination of the School of Nursing of Universidade Federal de Minas Gerais (*Observatório em Pesquisa e Estudo em Vacinação da Escola de Enfermagem da Universidade Federal de Minas Gerais* – OPESV-EEUFMG) in collaboration with the State Health Department of Minas Gerais (*Secretaria Estadual de Saúde do Estado de Minas Gerais* – SES-MG) was to evaluate the impact of the “Strategies for Increasing Vaccination Coverage in Children Under Two Years of Age in the State of Minas Gerais, Brazil: An Action Research” project on vaccination coverage among children, through execution of workshops and preparation of Action Plans individualized to the reality of each municipality in the state^{13,14}. This evaluation aimed to assess changes in vaccination coverage between the years 2021 and 2022, taking into account the size of the municipalities in the state.

METHODS

The study conducted within the state action research project was designed as a community clinical trial of the before-after type. It was carried out in 212 municipalities belonging to 08 Regional Health Managements and Superintendencies (*Gerências e Superintendências Regionais de Saúde* – GRS/SRS) in the state of Minas Gerais, Brazil, designed with the aim of organizing and managing health in the regions of the state¹⁴. All information was extracted from the National Immunization Program Information System (SIPNI), from 2021 and 2022, available at: <sipni.datasus.gov.br>.

The study encompassed 212 municipalities in Minas Gerais, selected based on their low vaccination coverage and a trend of decreasing coverage observed from 2015 to 2021¹². Workshops were conducted in collaboration with professionals responsible for vaccination and management in each municipality, aiming to develop and monitor action plans tailored to the specific context of each location. The health regions covered and the number of municipalities in each of them were: Regional Health Management of São João del Rey, made up of 20 municipalities; Alfenas Regional Health Superintendency, made up of 24 municipalities; Passos Regional Health Superintendency, made up of 27 municipalities; Leopoldina Regional Health Management, made up of 15 municipalities; Barbacena Regional Health Superintendency, made up of 31 municipalities; Ituiutaba Regional Health Management, made up of 9 municipalities; Regional Health Superintendency of Coronel Fabriciano, made up of 35 municipalities and Regional Health Superintendency of Governador Valadares, made up of 51 municipalities¹⁴.

The variables listed in Chart 1 were employed in this study. The 212 municipalities under evaluation were categorized according to the classification by Braz et al.¹⁰, which relies on criteria outlined in the indicators booklet of the Health Surveillance Actions Qualification Program (*Programa de Qualificação das Ações de Vigilância em Saúde* – PQAVS). Namely:

- Small municipality: One with a population of up to 20,000 inhabitants;
- Medium-sized municipality: One with a population between 20,001 and 100,000 inhabitants; and
- Large municipality: One with more than 100,000 inhabitants.

In this study, secondary data on Vaccination Coverage (VC), Homogeneity of Vaccination Coverage (HVC), and Abandonment rate of multi-dose vaccines for 12 immunobiologicals recommended for children under two years of age were utilized. The data spanned from January to December 2021 (pre-intervention period, prior to the state action research project) and January to December 2022 (post-intervention period).

The 12 analyzed immunobiologicals recommended for children under two years of age were: BCG, Hepatitis B (for children under one month of age), oral vaccine against rotavirus (second dose of Rotavirus vaccine in SUS and the second dose of Pentavalent Rotavirus in the private network), Meningococcal C (second dose of Meningococcal C and second dose of Meningococcal ACWY), Pneumococcal 10 and 13 (two doses), Pentavalent (third dose of the Pentavalent vaccine and third dose of the Hexavalent vaccine in the private network), Polio vaccine (third dose of Pentavalent and Hexavalent VIP/VOP in the private network), Triple Viral (two doses: second dose of Quadruple Viral, second dose, and single dose of Tetraviral), Yellow Fever (double dose, initial and first dose), Hepatitis A (considered

one dose), and Chickenpox (1 dose of Chickenpox and 1 dose of Tetraviral).

For VC, the classification of vaccination coverage established by PNI was adopted, according to which coverage must be greater than or equal to 90% for the BCG and oral Human Rotavirus vaccines and greater than or equal to 95% for the other immunizers. Vaccines are considered to have coverage very low if there is a VC between 0 and 50%; low if it is in the range of 50% to any value below the target; and adequate if it is greater than or equal to the target¹⁰.

HVC, an estimate of the proportion of municipalities with adequate coverage, agreed upon by the Organizational Contract for Public Health Action (*Contrato Organizativo de Ação Pública da Saúde* – COAP)¹⁰, is considered adequate when the municipality presents: from 75 to 100% of adequate coverage of at least ten vaccines; low when it presents between 50 and less than 75% adequate coverage of at least ten vaccines; and very low when the percentage reaches less than 50%¹⁰.

The abandonment rate of multi-dose vaccines, which contain multiple doses (such as the following vaccines: Pneumococcal 10 and 13, oral Human Rotavirus vaccine, and VIP/VOP Hexa and Pentavalent against Poliomyelitis), was also explored to understand the scenario of 2021 and 2022 and changes in relation to the abandonment of vaccination in municipalities through classification. It was considered low AP when less than 5% of people (<5%) stopped taking other doses; average when dropout varied from 5% to 10% (≥ 5 to <10%); and high when it reached a proportion greater than 10%.

Finally, the risk of the return of these diseases in these same municipalities was observed using the classification made by Braz et al.¹⁰, as follows:

- Very low risk: municipalities with 100% homogeneity;

Chart 1. Summary of the parameters used to calculate the immunization indicators in this study.

Indicator	Parameters
Vaccination Coverage Index (VCI)	Very low: <50% Low: $\geq 50\%$ and lower than target Adequate: \geq target
Homogeneity of Vaccination Coverage (HVC)	Adequate: $\geq 75\%$ to $\leq 100\%$ Low: $\geq 50\%$ to <75% Very low: <50%
Abandonment rate of multi-dose vaccines	Low: <5% Medium: $\geq 5\%$ to <10% High: $\geq 10\%$
Population size	Small size: Population $\leq 20,000$ inhabitants Medium size: Population between 20,001 and 100,000 inhabitants Large size: Population $\geq 100,001$ inhabitants
Risk Classification of Transmission of Vaccine-Preventable Diseases	Very low: municipalities with HVC =100% Low: municipalities with HVC $\geq 75\%$ and <100%, with adequate VCI for Polio, MMR and Pentavalent vaccines Medium: municipality with HVC $\geq 75\%$ and <100% and VCI below the target for one or more of the Polio, MMR or Pentavalent vaccines High: municipalities with HVC <75%, regardless of vaccination coverage Very high: municipality with HVC <75%, high TA ($\geq 10\%$) for any of the vaccines evaluated and with a large population size and, also, municipalities without vaccination records for any vaccine, regardless of population size

- Low risk: when there is adequate homogeneity (≥ 75 and $< 100\%$) of Polio, MMR and Pentavalent vaccines;
- Medium risk: ≥ 75 and 100% of homogeneity and adequate vaccination coverage for one of the three indicated vaccines;
- High risk: homogeneity $< 75\%$, regardless of indicator vaccines;
- Very high risk: homogeneity $< 75\%$ and proportion greater than 10% of abandonment rate of multi-dose vaccines evaluated in large municipalities, in places that do not have vaccination records, regardless of the number of inhabitants

Due to the low number of municipalities that present medium and very high risk, according to the proposed classification, in this study the categories were combined into: "low and medium risk" and "high and very high risk."

The analyses were conducted using the Statistical Software for Professionals (Stata), version 16.0. The aim was to understand the proportion of municipalities with low vaccination coverage for each of the 12 vaccine-preventable diseases analyzed, the proportion of homogeneity, and the percentage risk of vaccine-preventable diseases returning, depending on the size of municipalities. Additionally, the McNemar test was applied to verify the proportions of changes and their relevance, with a significance level set at 5%.

RESULTS

Following the implementation of the action research project, there was a notable increase in the proportion of municipalities classified as small that met the vaccination coverage targets established by the PNI. Significant increases ($p < 0.05$) were observed for several immunobiologicals: BCG (25.71% to 42.37%), Rotavirus (45.76% to 57.63%), Meningococcus (35.59% to 52.54%), Pneumococcal C (38.98% to 53.11%), Pentavalent (37.85% to 52.54%), Poliomyelitis (38.98% to 53.11%), MMR D2 (22.86% to 35.80%), and Chickenpox (40.11% to 59.89%), as detailed in Table 1.

For medium-sized municipalities, there was an increase in the number of municipalities reaching the target vaccination coverage after the survey, notably for BCG and Hepatitis B (first dose), with statistically significant differences observed. Conversely, among municipalities classified as large, there was no statistically significant increase in achieving the vaccination coverage target among children under two years of age (Table 1).

Tables 2, 3, and 4 illustrate the proportions of abandonment rate of multi-dose vaccines, HVC, and risk classification based on the population size of the municipalities. Significantly, among small municipalities, there was an increase in the proportion of those classified as having low abandonment rate of multi-dose vaccines for the Rotavirus vaccine (63.28% to 80.79%) after the implementa-

tion of the action research project. Moreover, there was an increase in the number of municipalities categorized with adequate homogeneity in vaccination coverage. Additionally, there was an increase in the number of municipalities classified as very low risk, low risk, and medium risk, coupled with a decrease in those classified as high risk and very high risk, all demonstrating statistically significant differences (Table 2).

No statistically significant difference was observed between municipalities classified as medium and large for any of the indicators evaluated in this study (Tables 3 and 4).

DISCUSSION

This study reaffirmed the importance of the activities promoted by the action research project, especially in small municipalities across all regions, while also noting the maintenance of the risk status in some municipalities. Rare exceptions showed a worsening in vaccination coverage. In light of these findings, it was pertinent to analyze the aspects that contributed to this outcome, with the aim of applying them in municipalities that have not yet reached the appropriate vaccination levels.

The extensive connection between primary health care and small municipalities may contribute to the effectiveness of actions in these cities¹⁵, given that approximately 177 of the municipalities studied have fewer than 20,000 inhabitants. According to the National Primary Care Policy (*Política Nacional de Atenção Básica - PNAB*)¹⁶, in areas characterized by significant territorial dispersion, risk, and social vulnerability, it is recommended to achieve 100% coverage of the population with a maximum of 750 people per Community Health Worker (CHW).

Moreover, CHWs play an extremely crucial role in transforming this scenario, engaging in health education through both individual and collective activities, and serving as a vital link between the professional team and the community¹⁶. With their intimate knowledge of the territory and its residents, CHWs can tailor interventions more effectively¹⁷. Training these CHWs to actively seek out and engage with individuals who are hesitant about vaccination, thus encouraging vaccination uptake, is essential. This is because CHWs are already familiar with local needs and have earned the trust of community members¹⁸. Therefore, in smaller municipalities, communication becomes more manageable as distances are shorter, and the number of individuals to be reached and informed is lower.

Small municipalities possess highly advantageous characteristics in line with the operational principles of SUS, particularly regionalization, which entails the decentralization of health services and actions tailored to the specific needs of each area, reflecting its historical and cultural context¹⁹. Consequently, addressing the health-care needs of small towns becomes more achievable as

Table 1. Percentage of municipalities according to vaccination coverage classification, Minas Gerais, 2021 and 2022.

	2021	2022	p-value
	% of municipalities that reached the target	% of municipalities that reached the target	
Small-sized municipalities (n=177)			
BCG	25.71	42.37	<0.001
Hepatitis B	17.51	23.16	0.131
Rotavirus	45.76	57.63	0.021
Meningococcal C	35.59	52.54	0.001
Pneumococcal C	38.98	53.11	0.008
Penta (DTP/Hib/HB)	37.85	52.54	0.006
Polio	38.98	53.11	0.008
MMR D1	46.33	55.93	0.077
Yellow Fever	33.33	34.46	0.816
MMR D2	22.86	35.80	0.002
Hepatitis A	42.94	50.85	0.126
Varicella	40.11	59.89	<0.001
Medium-sized municipalities (n=28)			
BCG	7.14	39.29	0.002
Hepatitis B	3.57	17.86	0.045
Rotavirus	17.86	28.57	0.256
Meningococcal C	7.14	14.29	0.317
Pneumococcal C	10.71	25.00	0.102
Penta (DTP/Hib/HB)	2.14	14.29	0.317
Polio	7.14	14.29	0.317
MMR D1	14.29	17.86	0.654
Yellow Fever	7.14	7.14	1.000
MMR D2	3.57	10.71	0.317
Hepatitis A	3.57	14.29	0.179
Varicella	3.57	21.43	0.058
Large-sized municipalities (n=7)			
BCG	14.29	57.14	0.083
Hepatitis B	14.29	0	0.317
Rotavirus	0	14.29	0.317
Meningococcal C	0	28.57	0.157
Pneumococcal C	0	28.57	0.157
Penta (DTP/Hib/HB)	0	28.57	0.157
Polio	0	28.57	0.157
MMR D1	0	14.29	0.317
Yellow Fever	0	0	-
MMR D2	0	0	-
Hepatitis A	0	14.19	0.317
Varicella	0	0	-

Note: 212 municipalities; target $\geq 90\%$ for the BCG and Human Rotavirus vaccine and $\geq 95\%$ for other immunobiologicals. 90% for the BCG and Oral Human Rotavirus vaccine and greater than or equal to 95% for other immunizers. Bold indicates statistically significant p-values.

it allows local authorities to have a deeper understanding of their community's dynamics. This aligns with the strategies employed in the action research, where each region tailored interventions based on identified weaknesses and strengths as highlighted by municipal managers. These interventions included securing funding, extending vaccination service hours, conducting outreach campaigns, and organizing professional development activities. Hence, the physical proximity inherent in small municipalities

likely played a pivotal role in the observed improvements within a single year²⁰.

The efficacy of interventions in medium and large municipalities also had a notable impact on vaccination coverage across Minas Gerais. Despite these advancements, sustaining progress in certain areas remains a challenge for further enhancing vaccination coverage among children under two years old and addressing apprehensions regarding the resurgence of vaccine-preventable diseases.

Table 2. Proportion of abandonment, homogeneity of vaccines and risk classification for the transmission of vaccine-preventable diseases, municipalities with a small population, Minas Gerais, 2021 and 2022.

Small-sized municipalities	2021	2022	p-value
	% of municipalities	% of municipalities	
Abandonment rate of multi-dose vaccines			
Human Rotavirus Oral Vaccine			
Low (<5%)	63.28	80.79	<0.001
Medium (≥5 to ≤10%)	17.51	11.30	
High (>10%)	19.21	7.91	
Pneumococcal 10 and 13			
Low (<5%)	64.97	71.75	0.174
Medium (≥5 to ≤10%)	15.25	14.69	
High (>10%)	19.77	13.56	
Pentavalent and Hexavalent			
Low (<5%)	62.71	60.45	0.679
Medium (≥5 to ≤10%)	11.86	18.08	
High (>10%)	25.42	21.47	
IPV/OPV, Hexavalent, and Pentavalent			
Low (<5%)	58.76	60.45	0.750
Medium (≥5 to ≤10%)	13.56	16.38	
High (>10%)	27.68	23.16	
Homogeneity of vaccination coverage			
Adequate for COAP (≥75 to <100%)	23.16	36.16	0.005
Low (≥50 to <75%)	18.64	18.08	
Very low (≥0 to <50%)	58.19	45.76	
Risk classification			
Very low risk	10.73	17.51	0.039
Low and medium risk	12.43	18.64	
High and very high risk	76.84	63.84	

Bold indicates statistically significant values.

Table 3. Proportion of abandonment, homogeneity of vaccines and risk classification for the transmission of vaccine-preventable diseases, municipalities with a medium population size, Minas Gerais, 2021 and 2022.

Medium-sized municipalities	2021	2022	p-value
	% of municipalities	% of municipalities	
Abandonment rate of multi-dose vaccines			
Human Rotavirus Oral Vaccine			
Low (<5%)	85.71	92.86	0.414
Medium (≥5 to ≤10%)	14.29	7.14	
High (>10%)	-	-	
Pneumococcal 10 and 13			
Low (<5%)	82.14	92.86	0.256
Medium (≥5 to ≤10%)	17.86	7.14	
High (>10%)	-	-	
Pentavalent and Hexavalent			
Low (<5%)	75.00	64.29	0.365
Medium (≥5 to ≤10%)	25.00	21.43	
High (>10%)	-	14.29	
IPV/OPV, Hexavalent, and Pentavalent			
Low (<5%)	71.43	64.29	0.527
Medium (≥5 to ≤10%)	25.00	25.00	
High (>10%)	3.57	10.71	
Homogeneity of vaccination coverage			
Adequate for COAP (≥75 to <100%)	-	10.71	0.083
Low (≥50 to <75%)	10.71	3.57	
Very low (≥0 to <50%)	89.29	85.71	
Risk classification			
Very low risk	-	-	-
Low and medium risk	-	10.71	
High and very high risk	100	89.29	

Table 4. Proportion of abandonment, homogeneity of vaccines and risk classification for the transmission of vaccine-preventable diseases, municipalities with a large population, Minas Gerais, 2021 and 2022.

Large-sized municipalities	2021	2022	p-value
	% of municipalities	% of municipalities	
Abandonment rate of multi-dose vaccines			
Human Rotavirus Oral Vaccine			
Low (<5%)	85.71	85.71	-
Medium (≥5 to ≤10%)	14.29	14.29	
High (>10%)	-	-	
Pneumococcal 10 and 13			
Low (<5%)	85.71	71.43	0.563
Medium (≥5 to ≤10%)	14.29	28.57	
High (>10%)	-	-	
Pentavalent and Hexavalent			
Low (<5%)	57.14	57.14	1.000
Medium (≥5 to ≤10%)	14.29	14.29	
High (>10%)	28.57	28.57	
IPV/OPV, Hexavalent, and Pentavalent			
Low (<5%)	57.14	57.14	1.000
Medium (≥5 to ≤10%)	14.29	14.29	
High (>10%)	28.57	28.57	
Homogeneity of vaccination coverage			
Adequate for COAP (≥75 to <100%)	-	-	-
Low (≥50 to <75%)	-	14.29	
Very low (≥0 to <50%)	100	85.71	
Risk classification			
Very low risk	-	-	-
Low and medium risk	-	-	
High and very high risk	100	100	

Limitations were observed concerning medium and large municipalities, as the improvements achieved were significant but still inadequate to decrease the percentages, unlike what was observed in small municipalities. This underscores weaknesses in the plans devised by regional authorities for these municipalities or in their implementation. Therefore, further investigations are necessary to gain a better understanding of the situation in these locations, as positive results were also observed in some municipalities within each of these regions.

In contrast to small municipalities, medium and large municipalities face challenges related to high population density. The population volume appears to be a complicating element in that it prevents the link between population and service, since, due to there being a large flow, health units tend to be unable to manage the quantity and possible issues that are obstacles in relation to vaccination, resulting in a failure in the responsibility of primary care in this aspect²¹. This burden on health professionals and CHWs affects the implementation of crucial actions like active search, which is more feasible in municipalities with smaller populations. Therefore, the failure to address these challenges and communicate them effectively to relevant authorities perpetuates the decline in vaccination coverage. It is essential to tailor interventions to the spe-

cific needs of each location to address territorial demands effectively and achieve positive outcomes across the state.

As vaccination coverage increased, municipalities also improved their risk status, accompanied by a decrease in the proportion of abandonment of multi-dose immunobiologicals. With more individuals vaccinated, the likelihood of diseases resurfacing diminishes²¹, and the awareness fostered — which contributed to the uptick in vaccination rates — may be correlated with the decline in abandonment of multi-dose vaccines.

It is important to acknowledge that this study has certain limitations. One such limitation is the reliance on data provided by SIPNI, which may be subject to underestimation due to potential shortcomings in the system's record-keeping.

Nevertheless, the methodological rigor employed in this study underscores the significance of a tailored approach to each municipality, with the goal of amplifying the positive outcomes observed in vaccination coverage improvement across Minas Gerais. Additionally, it is imperative for regional authorities to draw insights from the successful management strategies implemented in small municipalities and develop effective plans tailored to the unique characteristics of each locality, particularly those with persistently low vaccination coverage rates.

REFERENCES

1. Domingues CMAS, Maranhão AGK, Teixeira AM, Fantinato FFS, Domingues RAS. 46 anos do Programa Nacional de Imunizações: uma história repleta de conquistas e desafios a serem superados. *Cad Saúde Pública* 2020; 36 Supl 2: 00222919. <https://doi.org/10.1590/0102-311X00222919>
2. Domingues CMAS, Teixeira AMS. Coberturas vacinais e doenças imunopreveníveis no Brasil no período 1982-2012: avanços e desafios do Programa Nacional de Imunizações. *Epidemiol Serv Saúde* 2013; 22(1): 9-27. <https://doi.org/10.5123/S1679-49742013000100002>
3. Conselho Federal de Enfermagem. Taxa de vacinação infantil cai e o Brasil volta a patamar de 1987 [Internet]. 2022 [cited on May 14, 2023]. Available at: http://www.cofen.gov.br/taxa-de-vacinacao-infantil-cai-e-brasil-volta-a-patamar-de-1987_98834.html#:~:text=Queda%20na%20imuniza%C3%A7%C3%A3o%20das%20crian%C3%A7as,pior%20n%C3%ADvel%20em%20tr%C3%AAs%20d%C3%A9cadas
4. Sato APS. What is the importance of vaccine hesitancy in the drop of vaccination coverage in Brazil? *Rev Saude Publica* 2018; 52: 96. <https://doi.org/10.11606/S1518-8787.2018052001199>
5. Sato APS. Pandemia e coberturas vacinais: desafios para o retorno às escolas. *Rev Saude Publica* 2020; 54: 115. <https://doi.org/10.11606/s1518-8787.2020054003142>
6. Silva TMR, Sá ACMGN, Prates EJS, Saldanha RF, Silva TPR, Teixeira AMS, et al. Temporal and spatial distribution trends of polio vaccine coverage in less than one-year old children in Brazil, 2011-2021. *BMC Public Health* 2023; 23(1): 1359. <https://doi.org/10.1186/s12889-023-16192-8>
7. Arroyo LH, Ramos ACV, Yamamura M, Weiller TH, Crispim JA, Cartagena-Ramos D, et al. Areas with declining vaccination coverage for BCG, poliomyelitis, and MMR in Brazil (2006-2016): maps of regional heterogeneity. *Cad Saude Publica* 2020; 36(4): e00015619. <https://doi.org/10.1590/0102-311X00015619>
8. MacDonald NE; SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: definition, scope and determinants. *Vaccine* 2015; 33(34): 4161-4. <https://doi.org/10.1016/j.vaccine.2015.04.036>
9. Frugoli AG, Prado RS, Silva TMR, Matozinhos FP, Trapé CA, Lachtim SAF. Vaccine fake news: an analysis under the World Health Organization's 3Cs model. *Rev Esc Enferm USP* 2021; 55: e03736. <https://doi.org/10.1590/S1980-220X2020028303736>
10. Braz RM, Domingues CMAS, Teixeira AMS, Luna EJA. Classificação de risco de transmissão de doenças imunopreveníveis a partir de indicadores de coberturas vacinais nos municípios brasileiros. *Epidemiol Serv Saúde* 2016; 25(4): 745-54. <https://doi.org/10.5123/S1679-49742016000400008>
11. Silva TPR, Vimieiro AM, Gusmão JD, Souza JFA, Lachtim SAF, Vieira EWR, et al. Classificação de risco para transmissão de doenças imunopreveníveis em Minas Gerais, Brasil: dois anos desde o início da pandemia de COVID-19. *Ciênc Saúde Colet* 2023; 28(3): 699-710. <https://doi.org/10.1590/1413-81232023283.11192022>
12. Souza JFA, Silva TPR, Silva TMR, Amaral CD, Ribeiro EEN, Vimieiro AM, et al. Cobertura vacinal em crianças menores de um ano no estado de Minas Gerais, Brasil. *Ciênc Saúde Colet* 2022; 27(9): 3659-67. <https://doi.org/10.1590/1413-81232022279.07302022>
13. Souza JFA, Silva TPR, Lachtim SAF, Vieira EWR, Martins EF, Araújo DG, et al. The impact of strategies for increasing vaccination coverage in children: a community clinical trial. *Vaccine X* 2024; 16: 100429. <https://doi.org/10.1016/j.jvax.2024.100429>
14. Lachtim SAF, Palhoni ARG, Silva TPR, Ribeiro EEN, Souza JFA, Coelho VMR, et al. Estratégias cooperativas para melhorar a cobertura vacinal em crianças no estado de Minas Gerais, Brasil. *Arq Ciências Saúde UNIPAR* 2023; 27(9): 5310-23. <https://doi.org/10.25110/arqsaude.v27i9.2023-026>
15. Souza PA, Gandra B, Chaves ACC. Experiências sobre imunização e o papel da atenção primária à saúde. *APS em Revista* 2020; 2(3): 267-71. <https://doi.org/10.14295/aps.v2i3.57>
16. Brasil. Ministério da Saúde. Gabinete do Ministro. Portaria nº 2.436, de 21 de setembro de 2017. Aprova a Política Nacional de Atenção Básica, estabelecendo a revisão de diretrizes para a organização da Atenção Básica, no âmbito do Sistema Único de Saúde (SUS) [Internet]. Brasília: Ministério da Saúde; 2017 [cited on Feb 10, 2024]. Available at: https://bvsms.saude.gov.br/bvs/saudelegis/gm/2017/prt2436_22_09_2017.html
17. Costa SM, Araújo FF, Martins LV, Nobre LLR, Araújo FM, Rodrigues CAQ. Agente Comunitário de Saúde: elemento nuclear das ações em saúde. *Ciênc Saúde Colet* 2013; 18(7): 2147-56. <https://doi.org/10.1590/S1413-81232013000700030>
18. Costa GA, Alves LSF, Lima Júnior FO. A territorialização na Atenção Básica e o pensar estratégico na prevenção e promoção à saúde. *Revista Política e Planejamento Regional* 2022; 9(3): 395-407.
19. Carvalho ALB, Jesus WLA, Senra IMVB. Regionalização no SUS: processo de implementação, desafios e perspectivas na visão crítica de gestores do sistema. *Ciênc Saúde Colet* 2017; 22(4): 1155-64. <https://doi.org/10.1590/1413-81232017224.30252016>
20. Nassaralla APA, Doumit AM, Melo CF, Léon LC, Vidal RAR, Moura LR. Dimensões e consequências do movimento antivacina na realidade brasileira. *Rev Educ Saúde* 2019; 7(Supl. 1): 120-5.
21. Giovanella L, Bousquat A, Schenkman S, Almeida PF, Sardinha LMV, Vieira MLFP. Cobertura da Estratégia Saúde da Família no Brasil: o que nos mostram as Pesquisas Nacionais de Saúde 2013 e 2019. *Ciênc Saúde Colet* 2021; 26(Supl. 1): 2543-56. <https://doi.org/10.1590/1413-81232021266.1.43952020>
22. Brasil. Ministério da Saúde. Sistema de Informações do Programa Nacional de Imunizações [Internet]. DATASUS; 2022 [cited on Mar 5, 2024]. Available at: <https://sipni.datasus.gov.br/si-pni-web/faces/inicio.jsf>

RESUMO

Objetivo: Avaliar o impacto do projeto estadual de pesquisa-ação nas coberturas vacinais em crianças menores de dois anos no estado de Minas Gerais, segundo o porte dos municípios, comparando os anos de 2021 e 2022. **Métodos:** Estudo aninhado ao projeto estadual de pesquisa-ação, um ensaio clínico comunitário, do tipo antes-depois, realizado em 212 municípios do estado de Minas Gerais. Utilizou-se, neste estudo, dados secundários de Cobertura Vacinal (CV), Homogeneidade das Vacinas (HCV) e Proporção de Abandono (PA). Após a classificação de municípios por porte, e as taxas de cobertura vacinal serem classificadas de maneira equânime, foi realizada uma análise de dados secundários de 12 imunobiológicos indicados para a faixa etária em questão e de suas PAs. **Resultados:** Observou-se aumento na proporção de municípios classificados como de pequeno porte que atingiram a meta da cobertura vacinal estabelecida pelo Programa Nacional de Imunização (PNI), após a realização do projeto de pesquisa-ação. Houve aumento na proporção de municípios de pequeno porte classificados como de baixa PA para a vacina Rotavírus, na homogeneidade adequada da CV e na classificação de risco para risco muito baixo e risco baixo e médio, todos com diferença estatisticamente significativa. **Conclusão:** Houve influência do porte municipal na efetividade das ações aplicadas para aumento de cobertura vacinal, explicitando que a proposição de ações individualizadas para cada município é imprescindível para melhorar a cobertura vacinal.

Palavras-chave: Vacinação. Cobertura vacinal. Doenças transmissíveis. Municípios. Epidemiologia.

AUTHORS' CONTRIBUTIONS: Moreira, C.M.: Formal analysis, Conceptualization, Data curation, Writing – original draft, Writing – review & editing, Methodology, Validation. Silva, T.P.R.: Formal analysis, Conceptualization, Data curation, Writing – original draft, Writing – review & editing, Methodology, Validation. Neves, M.C.A.: Formal analysis, Conceptualization, Data curation, Writing – original draft, Writing – review & editing, Methodology, Validation. Cruz, M.V.G.: Conceptualization, Data curation, Writing – review & editing. Ribeiro, E.E.N.: Writing – review & editing. Ferreira Júnior, S.: Writing – review & editing. Lachtim, S.A.F.: Writing – review & editing. Matozinhos, F.P.: Formal analysis, Conceptualization, Data curation, Writing – review & editing, Methodology, Writing – original draft, Validation, Supervision.

FUNDING: none.

ACKNOWLEDGMENTS: To the Vaccination Research and Studies Observatory of the Nursing School of Universidade Federal de Minas Gerais (*Observatório de Pesquisa e Estudos em Vacinação da Escola de Enfermagem da Universidade Federal de Minas Gerais – OPESV*) and the State Department of Health of Minas Gerais (*Secretaria de Estado de Saúde de Minas Gerais – SES-MG*) for their support in conducting this study.



© 2024 | *Epidemiologia* is a publication of

Associação Brasileira de Saúde Coletiva - ABRASCO